

ESD-TR-66-66

MTR-32

# ESD RECORD COPY

RETURN TO  
SCIENTIFIC & TECHNICAL INFORMATION DIVISION  
(ESTI), BUILDING 1211

## ESD ACCESSION LIST

ESTI Call No. AL 51345  
Copy No. 1 of 1 cys.

### NUMERICAL COMPARISON OF EXACT, UNIFORM, AND NON-UNIFORM ASYMPTOTIC SOLUTION OF THE REDUCED WAVE EQUATION NEAR A CAUSTIC

JUNE 1966

R. M. Lewis (Consultant)

L. J. Kaplan

Prepared for

SPACE DEFENSE SYSTEM PROGRAM OFFICE (496L/474L)

ELECTRONIC SYSTEMS DIVISION  
AIR FORCE SYSTEMS COMMAND  
UNITED STATES AIR FORCE  
L. G. Hanscom Field, Bedford, Massachusetts



Distribution of this document is unlimited.

Project 4966

Prepared by

THE MITRE CORPORATION

Bedford, Massachusetts

Contract AF19(628)-5165

55X

ADD 635049

This document may be reproduced to satisfy official needs of U.S. Government agencies. No other reproduction authorized except with permission of Hq. Electronic Systems Division, ATTN: ESTI.

When US Government drawings, specifications, or other data are used for any purpose other than a definitely related government procurement operation, the government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Do not return this copy. Retain or destroy.

NUMERICAL COMPARISON OF EXACT, UNIFORM, AND NON-UNIFORM  
ASYMPTOTIC SOLUTION OF THE REDUCED WAVE EQUATION NEAR A CAUSTIC

JUNE 1966

R. M. Lewis (Consultant)

L. J. Kaplan

Prepared for

SPACE DEFENSE SYSTEM PROGRAM OFFICE (496L/474L)

ELECTRONIC SYSTEMS DIVISION  
AIR FORCE SYSTEMS COMMAND  
UNITED STATES AIR FORCE  
L. G. Hanscom Field, Bedford, Massachusetts



Distribution of this document is unlimited.

**Project 4966**

Prepared by

THE MITRE CORPORATION  
Bedford, Massachusetts

Contract AF19(628)-5165

## FOREWORD

We wish to thank D. Ludwig and J. D. R. Kramer for their interesting comments and suggestions. We also wish to thank Mrs. M. J. Carey for her able programming.

## REVIEW AND APPROVAL

This technical report has been reviewed and is approved.



HARRY R. BETZ, 2/Lt., USAF  
Project Manager  
496L/474L System Program Office

## ABSTRACT

A numerical comparison of the exact solution, the uniform asymptotic solution, and the non-uniform asymptotic solution of a field that produces a circular caustic is performed. It is seen that the uniform asymptotic solution is accurate even at the caustic for moderate values of  $ka$ . Curves and data are given as a function of  $ka$  and distances from the caustic. The asymptotic expressions are derived both from the exact solution and from the physics of the circular caustic.

## TABLE OF CONTENTS

	<u>Page</u>
SECTION I      DERIVATION	1
SECTION II     NUMERICAL RESULTS	7
APPENDIX I    THE CIRCULAR CAUSTIC	11
APPENDIX II   THE AIRY FUNCTION	16
APPENDIX III  THE UNIFORM ASYMPTOTIC FIELD AT THE CAUSTIC	18

## LIST OF ILLUSTRATIONS

### Figure Number

1	Circular Caustic Corresponding to Equation (15)	4
2	Plot of Field versus Normalized Distance for Various $ka$	7
3	Plot of Error versus Relative Distance for Two Values of $ka$	8
4	Plot of Field versus Normalized Distance for Various $ka$	8
5	Plot of Error on the Caustic as a Function of $ka$	10
6	Circular Caustic	12
7	Contour Relevant to the Airy Function	16

## SECTION I

### DERIVATION

The uniform asymptotic expansion developed in Lewis' paper\* is valid in the neighborhood of a caustic, and it reduces to the usual geometrical optics representation away from the caustic. In Appendix I, these expansions are calculated for the circular caustic. Herein, an exact solution of

$$\left(\nabla^2 + k^2\right) \mu = 0, \quad (1)$$

whose asymptotic expansions agree with that of Appendix I, enables us to obtain a numerical comparison of an exact solution and its uniform and non-uniform approximations.

An exact solution of (1) in two dimensions, where polar coordinates are used, is the function

$$\mu = e^{ika\phi} J_{ka}(kr). \quad (2)$$

Here  $J_n(z)$  is the Bessel function of order  $n$ . We obtain a uniform asymptotic expansion of (2) by using the following formula, given in Asymptotic Approximations by H. Jeffreys:<sup>†</sup>

$$J_n(n \operatorname{sech} \mu) \sim \left(\frac{4\xi}{\tan^2 \mu}\right)^{1/4} n^{-1/3} \operatorname{Ai}\left(n^{2/3} \xi\right); \quad \xi = \frac{3}{2} (\mu - \tanh \mu). \quad (3)$$

---

\* R. M. Lewis, Uniform Representation of Geometrical Optics Fields Near a Caustic, The MITRE Corporation, ESD-TR-65-404, Bedford, Mass., May 1966.

† H. Jeffreys, Asymptotic Approximations, England, Oxford University Press, 1962, p. 84, Equation (36).

In (3),  $Ai$  is the Airy function defined and discussed in Jeffreys' work\* and in Appendix II. The formula is valid for large  $n$ .

In order to apply (3) to (2), one sets

$$z - \operatorname{sech} \mu = (\cos v)^{-1}, \quad v = i\mu \quad (4)$$

Thus,

$$\frac{2}{3}\xi^{3/2} = -iv - \tanh(-iv) = i(\tan v - v) = i\left(\sqrt{z^2 - 1} - \cos^{-1} 1/z\right),$$

$$\tanh^2 \mu = 1 - \operatorname{sech}^2 \mu = 1 - z^2, \quad (5)$$

and

$$J_n(nz) \sim \left[ \frac{4\xi}{1 - z^2} \right]^{1/4} n^{-1/3} Ai\left(n^{2/3}\xi\right). \quad (6)$$

Here

$$-\xi = \frac{3}{2} \left[ \sqrt{z^2 - 1} - \cos^{-1} (1/z) \right]^{2/3} \quad (7)$$

in agreement with (5). Let  $n = ka$ ,  $z = r/a$  and apply (6) to (2). The result is

$$\mu \sim e^{ika\phi} \left[ \frac{4\xi}{1 - r^2/a^2} \right]^{1/4} (ka)^{-1/3} Ai\left[(ka)^{2/3}\xi\right]; \quad (8)$$

or

$$\mu \sim e^{ik\theta_g} Ai\left(-k^{2/3}\rho\right). \quad (9)$$

\*See page 1, second reference.



Here

$$\theta = a\phi, \quad (10)$$

$$\rho = -a^{2/3} = (3/2 \Psi)^{2/3}, \quad 2/3 \rho^{3/2} = \Psi = \sqrt{r^2 - a^2} - a \cos^{-1}(a/r), \quad (11)$$

and

$$g = (ka)^{-1/3} \left[ \frac{-4\xi}{r^2/a^2 - 1} \right]^{1/4} = \left[ \frac{4k^{2/3} \rho}{k^2(r^2 - a^2)} \right]^{1/4}. \quad (12)$$

The non-uniform expansion is obtained from (9) by using the asymptotic expansion

$$\text{Ai}(-\xi) \sim \frac{1}{\sqrt{\pi}} \xi^{-1/4} \sin\left(2/3 \xi^{3/2} + \pi/4\right); \quad -\pi/3 < \arg \xi < \pi/3; \quad |\xi| \rightarrow \infty \quad (13)$$

of the Airy function.\* Let

$$\xi = k^{2/3} \rho. \quad \text{Then } 2/3 \xi^{3/2} = k\Psi \text{ and}$$

$$\mu \sim \frac{1}{\sqrt{\pi}} e^{ika\phi} \left[ \frac{4}{k^2(r^2 - a^2)} \right]^{1/4} \sin(k\Psi + \pi/4), \quad (14)$$

or

$$\mu \sim \frac{1}{2\pi i} \left[ k^2(r^2 - a^2) \right]^{-1/4} \{ e^{ik(a\phi + \Psi) + i\pi/4} - e^{ik(a\phi - \Psi) - i\pi/4} \}. \quad (15)$$

This result may be interpreted in terms of geometrical optics with the aid of Figure 1.

Two rays, each tangent to the circular caustic of radius  $a$ , intersect at the point  $X$  whose polar coordinates are  $r, \phi$ . The two phases associated with these rays are

---

\* See page 1, second reference, p. 33, Equation (30).




Figure 1. Circular Caustic Corresponding to Equation (15)

$$S_{\pm} = \sigma_{\pm} \pm \tau, \quad (16)$$

where

$$\sigma_{\pm} = a(\phi \mp \alpha) = a\phi \mp a \cos^{-1} \left( \frac{a}{r} \right), \tau = \sqrt{r^2 - a^2}. \quad (17)$$

We see from (11) that

$$a\phi \pm \Psi = a\phi \mp a \cos^{-1} \left( \frac{a}{r} \right) \pm \tau = \sigma_{\pm} \pm \tau = S_{\pm}, \quad (18)$$

hence, (15) can be written in the form

$$\mu \sim \frac{e^{i\pi/4}}{\sqrt{2\pi k}} \left[ z_- e^{iks_-} + z_+ e^{iks_+ - i\pi/2} \right], \quad (19)$$

where

$$z_- = z_+ = \tau^{-1/2} . \quad (20)$$

In (20), the amplitude and the phase of the incoming and outgoing waves are clearly exhibited. The amplitudes become infinite at the caustic ( $\tau=0$ ), and the phase-shift factor  $e^{-i\pi/2}$  multiplies the outgoing wave. The wave fronts are the spirals  $S_{\pm} = \text{constant}$ .

Here, the uniform expansion (9-12) has been obtained directly from the exact solution (2). In general, of course, the exact solution is not known. Then, it is necessary to use the method developed in Lewis' report\* to obtain the uniform expansion. In Appendix I, that method is applied to this problem, and a uniform expansion is obtained that agrees exactly with (9-12).

Numerical computations and graphs of the exact, uniform, and non-uniform results will not be obtained. For this purpose,  $\phi=0$  may be taken, since  $e^{ika\phi}$  is a common factor of all three results. It is noted that  $ka$ ,  $kr$ ,  $k\Psi$ , and  $k^{2/3}\rho$  are dimensionless, therefore let  $k=1$ . The exact, uniform, and non-uniform results are denoted by  $\mu_1$ ,  $\mu_2$ , and  $\mu_3$ . Then,

$$\mu_1 = J_a(r), \quad (21)$$

$$\mu_2 = \left( \frac{4\rho}{r^2 - a^2} \right)^{1/4} \text{Ai}(-\rho), \quad (22)$$

and

$$\mu_3 = \sqrt{\frac{2}{\pi}} \left( \frac{1}{r^2 - a^2} \right)^{1/4} \sin(\Psi + \pi/4). \quad (23)$$

---

\* See page 1, first reference.

Here

$$\Psi = \sqrt{r^2 - a^2} - a \cos^{-1} (a/r), \quad (24)$$

and

$$\rho = \left( \frac{3}{2} \Psi \right)^{2/3}. \quad (25)$$

## SECTION II

### NUMERICAL RESULTS

These three functions were programmed for several values of  $ka$  as a function of  $r/a$ . Figure 2 shows these curves for  $ka = 2, 5$ , and  $10$ . For these values of  $ka$ , curves of the exact and uniform solutions are indistinguishable, and only curves of  $\mu_1$  and  $\mu_3$  are shown. To indicate the errors in the uniform approximations, a deviation curve is plotted for  $ka = 1$  and  $ka = 2$  in Figure 3. Figure 4 is the same as Figure 2 except that  $ka = 1/2, 1/4, 1/8$ , and  $1/16$ .

For these values of  $ka$ ,  $\mu_2$  has diverged sufficiently from  $\mu_1$  so that three curves can be distinguished. From these curves the following conclusions are clear. The non-uniform approximation is very accurate




Figure 2. Plot of Field versus Normalized Distance for Various  $ka$




Figure 3. Plot of Error versus Relative Distance for Two Values of  $ka$




Figure 4. Plot of Field versus Normalized Distance for Various  $ka$

for sufficiently large  $r/a$  and all  $ka$ . The uniform approximation is good for almost all  $r/a$  even for quite small  $ka$ . To indicate the actual errors in the uniform approximation, the maximum full-scale percentage error for various  $ka$  is indicated in Table I. This is defined as

$$\frac{\text{maximum } |\mu_2 - \mu_1| 100}{\text{maximum } \mu_1}.$$

From the curves of Figure 3, it can be seen that the deviation from the exact solution at a given point may be considerably smaller than this value.

Examination of the field on the caustic using (21), (22), and (23) leads to the exact solution  $\mu_1 = J_{ka}(ka)$ , while the uniform approximation is indeterminate and the non-uniform approximation is singular. In Appendix III, an expression is developed for the uniform approximation (26).

Table I  
Percentage Error for Various  $ka$

$ka$	Percentage Error
1/64	87.5
1/32	57.6
1/16	31.6
1/8	17.3
1/4	8.05
1/2	3.3
1	1.7
2	0.45
3	0.27
5	0.18
10	0.082
20	0.037
50	0.017



$$u = \frac{2^{1/3}}{a^{1/3}} A_i \left[ \frac{-2^{1/3} \epsilon}{a^{1/3}} \right] \quad (26)$$

In Figure 5, the actual percentage error at the caustic as a function of  $ka^*$  is plotted. Since the largest value of the exact solution is not at the caustic, these errors are slightly larger than those of Table I. As  $ka$  becomes large, the error goes to zero. As  $ka$  goes to zero,  $\mu_2$  becomes singular. At small  $ka$  the exact solution approaches unity, and hence the relative error,  $(\mu_2 - \mu_1)/\mu_1$ , approaches the difference.




Figure 5. Plot of Error on the Caustic as a Function of  $ka$

\* This curve was suggested by D. Ludwig, Courant Institute, New York, N. Y.



# APPENDIX I

## THE CIRCULAR CAUSTIC

Herein, aspects of the application of various representations of geometric optics fields near a caustic shall be considered. The standard asymptotic expansion and a representation due to R. M. Lewis\* shall also be considered. To be specific, the two-dimensional field that produces a circular caustic was chosen.

The wave front that produces a circular caustic is a spiral determined by a circle and constant length string. In Figure 6, we have the circular caustic of radius  $a$ . The points  $1'$  and  $2'$  lie on the spiral wave front. They are determined by the condition

$$\alpha_i + \rho_i = L, \quad (27)$$

where  $L$  is a constant and  $\rho_i$  is the distance on the wave front to the caustic; i. e.,  $\rho_i$  is the magnitude of the line segment  $11'$ .

The field at any point off the caustic is determined by the two rays that pass through the point. In Figure 2, the two indicated rays pass through the point  $P$ . The distance from the point  $i$  on the wave front to  $P$  is designated by  $\tau_i$ . The distances from  $P$  to 1 and 2 are equal and are

$$S = \sqrt{r^2 - a^2}, \quad (28)$$


where  $r \geq a$  is the radial coordinate of the point  $P$ . We note that

$$\tau_1 = \rho_1 - S, \quad (29)$$

$$\tau_2 = \rho_2 + S, \quad (30)$$

---

\* See page 1, first reference.



IA-16,779

Figure 6. Circular Caustic

or that one ray through  $\rho$  has touched the caustic. The angular location of points 1 and 2 are found from

$$\phi = \theta \pm \cos^{-1} a/r, \quad (31)$$

where  $\theta$  is the angular coordinates of P. Using (30), (27), and (28) on (29), the following is obtained:

$$\tau_1 = L - a \left( \theta - \cos^{-1} a/r \right) - \sqrt{r^2 - a^2}; \quad (32)$$

$$\tau_2 = L - a \left( \theta - \cos^{-1} a/r \right) + \sqrt{r^2 - a^2}. \quad (33)$$

The standard asymptotic formula\* for the field at  $\rho$  is

$$u = Z_1 \left( 1 - \tau_1/\rho_1 \right)^{-1/2} e^{ik(s_1 + \tau_1)} + Z_2 \left( 1 - \tau_2/\rho_2 \right)^{-1/2} e^{ik(s_2 + \tau_2)}, \quad (34)$$

where the subscript refers to the ray. This expression differs from that in Lewis' report\* since this is a two-dimensional problem where  $\rho$  is used instead of the curvature  $\kappa$ . The term associated with the second ray involves taking the square root of a negative quantity. This indicates the phase shift of  $\pi/2$  at the caustic. This formula is not valid at the caustic since it is singular at such points. Using the previous results for the circular caustic, relation (34) becomes

$$u = \frac{Z_1 (r^2 - a^2)^{-1/4}}{(L - a \cos^{-1} a/r)^{1/2}} e^{ik(s_1 + L - a\theta + a \cos^{-1} a/r - \sqrt{r^2 - a^2})} \\ + \frac{Z_2 (r^2 - a^2)^{-1/4}}{(1 - a \cos^{-1} a/r)^{1/2}} e^{ik(s_2 + L - a\theta - a \cos^{-1} a/r + \sqrt{r^2 - a^2} - \pi/2k)} \quad (35)$$

---

\* See page 1, first reference, Equation (93).

For computation, the following assumptions are convenient,

$$Z_1 e^{iks_1} = Z_2 e^{iks_2} = k^{1/2} (L - a \cos^{-1} a/r)^{1/2}, \quad (36)$$

$$\theta = 0. \quad (37)$$

Equation (37) requires that the field is constant along the wave front, and the results must be normalized so that this constant is unity. That  $\theta = 0$  follows from a reasonable choice of coordinate system.

To simplify the result,  $L$  can now be chosen such that

$$e^{ikL} e^{-i\pi/4} = 1. \quad (38)$$

Then (35) reduces to

$$u = 2i \frac{e^{ikL} e^{+i\pi/4}}{k^{1/2} (r^2 - a^2)^{1/4}} \sin \left( ka \cos^{-1} a/r - k \sqrt{r^2 - a^2} + \pi/4 \right), \quad (39)$$

and Equation (39) is rewritten

$$u = \frac{2}{k^{1/2} (r^2 - a^2)^{1/4}} \sin \left( ka \cos^{-1} a/r - k \sqrt{r^2 - a^2} + \pi/4 \right). \quad (40)$$

Lewis\* gives a uniform asymptotic expansion

$$u = \pi^{1/2} |\xi|^{1/2} e^{-i\pi/4} \left\{ Ai \left| -\xi^2 \right| \left[ \frac{Z_1 e^{ikb_1}}{|1 - \tau_1/\rho_1|^{1/2}} + \frac{Z_2 e^{ikb_2}}{|1 - \tau_2/\rho_2|^{1/2}} \right] \right. \\ \left. + \frac{i Ai' \left( -\xi^2 \right)}{\xi} \left[ \frac{Z_1 e^{ikb_1}}{|1 - \tau_1/\rho_2|^{1/2}} - \frac{Z_2 e^{ikb_2}}{|1 - \tau_2/\rho_2|^{1/2}} \right] \right\}. \quad (41)$$

---

\* See page 1, first reference, Equations (68) and (69).

where

$$\xi = (3k/4)^{1/3} (\tau_2 - \tau_1)^{1/3} \quad (42)$$

and

$$b_i = s_i + \frac{\tau_1 + \tau_2}{2} \quad (43)$$

Using Equations (32), (33), (36-38), and (41),

$$\mu = 2\pi^{1/2} \frac{\left| (3k/2) \left( \sqrt{r^2 - a^2} - a \cos^{-1} a/r \right) \right|^{1/6}}{k^{1/2} (r^2 - a^2)^{1/4}} \quad (44)$$

$$\text{Ai} \left( - \left[ (3k/2) \left| \sqrt{r^2 - a^2} - a \cos^{-1} a/r \right| \right]^{2/3} \right)$$

Note the disappearance of the derivative of the Airy function because of the symmetry of the caustics. If the asymptotic expansion of the Airy function\* is used in (44), Equation (40) is recovered.

The comparison of Equations (40) and (44) with Equations (22) and (23) shows agreement except for a numerical constant of  $\sqrt{1/2\pi}$ . Thus, a check is provided on the validity of the manipulations. This factor arises from the arbitrary choice of amplitude for the wave.

---

\* See page 1, second reference, p. 33, Equation (30).

## APPENDIX II

### THE AIRY FUNCTION

The Airy function is defined (e. g. , in Jeffreys' work)\* as

$$\text{Ai}(z) = \frac{1}{2\pi i} \int_{L_1} e^{tz - 1/3 t^3} dt , \quad (45)$$

where  $L_1$  is the contour in the  $t$ -plane shown in Figure 7.

The transformation  $t = is$  is introduced. Then for real  $z$  the transformed contour can be shifted to the real axis, and




Figure 7. Contour Relevant to the Airy Function

\*See page 1, second reference, Equation (1).

$$\text{Ai}(z) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(xz+s^3/3)} ds ; z \text{ real} \quad (46)$$

is obtained.

An alternative representation is

$$\text{Ai}(z) = \frac{1}{\pi} \int_0^{\infty} \cos(sz+s^3/3) ds ; z \text{ real} . \quad (47)$$

### APPENDIX III

#### THE UNIFORM ASYMPTOTIC FIELD AT THE CAUSTIC\*

A quick examination of Equation (44) or, equivalently, (22) does not indicate that the uniform asymptotic expansion is finite at the caustic. In Lewis' report<sup>†</sup>, it was shown that this was generally true. Here, this shall be shown directly from Equation (22). The resultant formula will also give a simple expression near the caustic. Let

$$r = a (1 + \epsilon) , \quad (48)$$

where  $\epsilon$  is a small positive quantity. Then

$$\sqrt{r^2 - a^2} = a \sqrt{2\epsilon + \epsilon^2} \sim a \sqrt{2\epsilon} , \quad (49)$$

and from relation (24)

$$\frac{\Psi - \sqrt{r^2 - a^2}}{a} = \cos^{-1} (a/r) . \quad (50)$$

Taking the cosine of both sides and expanding the power series to second order in  $\epsilon$  and first order in  $\Psi$  ,

$$\Psi = \frac{2\sqrt{2}}{3} \frac{\epsilon^{3/2}}{a^{1/2}} \quad (51)$$

is obtained. For the field near the caustic, the following can now be obtained.

$$u = \left(\frac{2}{a}\right)^{1/3} \text{Ai} \left[ -\epsilon \left(\frac{2}{a}\right)^{1/3} \right] . \quad (52)$$

---

\*We wish to thank J. D. R. Kramer of The MITRE Corporation, Bedford, Mass. for this derivation.

<sup>†</sup>See page 1, first reference, Equation (6).



DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) The MITRE Corporation Bedford, Massachusetts		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE Numerical Comparison of Exact, Uniform, and Non-uniform Asymptotic Solution of the Reduced Wave Equation Near a Caustic			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) N/A			
5. AUTHOR(S) (Last name, first name, initial) Lewis, R. M. (Consultant) Kaplan, L. J.			
6. REPORT DATE June 1966		7a. TOTAL NO. OF PAGES 20	7b. NO. OF REFS 2
8a. CONTRACT OR GRANT NO. AF19(628)-5165		9a. ORIGINATOR'S REPORT NUMBER(S) ESD-TR-66-66	
b. PROJECT NO. 4966			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		MTR-32	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Space Defense System Program Office (496L/474L); Electronic Systems Division, L. G. Hanscom Field, Bedford, Massachusetts	
13. ABSTRACT <p>A numerical comparison of the exact solution, the uniform asymptotic solution, and the non-uniform asymptotic solution of a field that produces a circular caustic is performed. It is seen that the uniform asymptotic solution is accurate even at the caustic for moderate values of <math>ka</math>. Curves and data are given as a function of <math>ka</math> and distances from the caustic. The asymptotic expressions are derived both from the exact solution and from the physics of the circular caustic.</p>			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
MATHEMATICS Caustic Geometrical Theory of Diffraction Asymptotic Expansions						

#### INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.